

IN COOPERATION WITH THE U.S. GEOLOGICAL SURVEY





Map	Name	Map coordinates	Development category	Resources*	Туре	Brief description	Principal reference
A	Stevens Creek	60 <sup>0</sup> 57′N, 157 <sup>0</sup> 21′W approx.	р	W	Vein	Wolframite in vein-quartz float from hornfelsed Kuskokwim Group sedimentary rocks.	Cobb, 1976, location 7, p. 76.
В	Taylor Creek	60°53′ - 60°54′N, 157°15′ - 157°22′W	M	Au (Hg, Sn)	Placer	Placer deposits contain gold, cinnabar, cassiterite, and pyrite. Production mainly in 1950-51, was about 2,500 ounces of fine gold. Placers were probably derived from contact zones around felsic bodies that hornfelsed clastic Cretaceous rocks.	Cobb, 1976, location 11, p. 77.

# Key: Development category

M - Mine with known production. Mine is defined as a mineral deposit with recorded production

p - Prospect work, but no known activity since 1960. Prospect is defined as a deposit that has been staked with subsequent exploratory or development work, but has no known production. \*Minor constituents or potential byproducts in parentheses.


## Table 6. Chemical data.

Map no.	30*	33+	49+
Sample no.	44ACa26	82MR309	83MR234
Lab. no.		15187	15239
Rock type	Porphyritic	Granite	Porphyritic
	granite		granite
		-	
Oxides		Percentage	
SiO <sub>2</sub>	70.48	69.24	68.50
$Al_2O_3$	13.97	14.12	15.41
$Fe_2O_3$	0.43	2.72	1.20
FeO	3.21	3.14	3.15
MgO	0.29	0.92	0.85
CaO	1.18	1.49	1.52
$Na_2O$	2.77	3.25	3.04
K <sub>2</sub> O	5.48	4.04	4.95
TiO <sub>2</sub>	0.45	0.57	0.54
P <sub>2</sub> O <sub>5</sub>	0.22	0.21	0.32
MnO	0.04	0.09	80.0
LOI	0.90	0.53	0.62
TOTAL	99.85	100.32	100.16
	CIPW norm	ative minerals	
Q	29.27	29.72	26.29
С	1.89	2.20	3.08
or	32.87	23.92	29.47
ab	23.79	27.56	25.92
an	4.48	6.03	5.49
hy	5.68	5.05	6.21
mg	0.63	3.95	1.75
il	0.87		1.03
ар	0.52	0.49	0.75
TOTAL	100.00	98.92	99.99

\*Analyses by USGS Chemistry and Physics laboratory, J.G. Fairchild analyst. +Analyses by X-ray fluorescence, DGGS Minerals Laboratory. LOI = Loss on ignition.

### INTRODUCTION This map is one in a series of geologic data bases (scale 1:63,360) from southwest Alaska. The series is scheduled for release by DGGS during 1985. Each map is a summary of geologic field data, analytical data, fossil reports, resource information, and land status available for that quadrangle. Geologic data were obtained primarily from U.S. Geological Survey (USGS) field notes collected from 1941 to 1975. Field locations were replotted on modern topographic base maps from the original trimetrogon air photos and reconnaissance topographic maps. The map-numbering sequence, which follows a normal written-page progression using the township and range mile-square blocks, starts in the northwest corner of the quadrangle and ends in the southeast corner. Where necessary, original station numbers were modified to fit the format: year, geologist, and field number (table 1). Initial station numbers commonly consisted of the date followed by a number for each observation point, starting each day with number 1, for example, 6/11-1. The modified number preserves the original number, but adds the year and geologist to eliminate confusion and ambiguity. For example, 47Hr6/11-1 was the first station occupied by Joseph M. Hoare on June 11, 1947. Rock descriptions (table 2) and structural data (table 3) were compiled from field notes exactly as written; no attempt was made to modify or interpret the original data. We appreciate the review by M.S. Robinson (DGGS) and Stephen E. Box (USGS).


# GENERALIZED DISTRIBUTION OF ROCK UNITS



# Platt, 1955; Blodgett and Clough, 1985) **EXPLANATION**

Quaternary deposits Granite, porphyritic granite, and dikes and sills of rhyolite Hornfelsed Kuskokwim Group sedimentary rocks Kuskokwim Group sedimentary rocks DSab - Early Devonian to Late Silurian

algal boundstone



State tentatively approved

### 0.786 $65.3 \pm 2.0$ x = 8.58049 83MR234C Porphyritic 7.370 7.296 x = 7.3439.110 0.3118 0.939 $65.0 \pm 2.0$ 9.200 $\bar{x} = 9.155$ 0.872 $64.5 \pm 1.9$ 83MR235 9.876

K-Ar age determinations by J.D. Blum and D.L. Turner, Alaska Division of Geological and Geophysical Surveys, University of Alaska,

## REFERENCES CITED

Blodgett, R.B., and Clough, J.B., 1985, The Nixon Fork terrane - Part of an in-situ peninsular extension of the North American Paleozoic continent: Geological Society of America, Abstracts with Programs [abs.] [in press].

kwim region, Alaska: U.S. Geological Survey Professional Paper 268, 132 p.

Cobb, E.H., 1976, Summary of references to mineral occurrences (other than mineral fuels and construction materials) in the Dillingham, Sleetmute, and Taylor Moun-

# COMPILATION OF GEOLOGIC DATA FROM THE TAYLOR MOUNTAINS D-4 QUADRANGLE, SOUTHWESTERN ALASKA

157°37 30

MAP SYMBOLS

mine or prospect (table 4)

extent of mine or prospect

78 - map number; corresponds

to table numbers

APPROXIMATE MEAN

**DECLINATION**, 1954

Compiled by

Rocky R. Reifenstuhl, John Decker, and Warren L. Coonrad

ALASKA

QUADRANGLE LOCATION

Base from U.S. Geological Survey

This report is a preliminary publica-

tion of DGGS. The authors are

solely responsible for its content

and will appreciate candid com-

ments on the accuracy of the data as well as suggestions to improve

Taylor Mountains D-4, 1954,

Quadrangle, Alaska

the report.

CONTOUR INTERVAL 50 FEET

DATUM MEAN SEA LEVEL

joint N63E, 53SE N77E, 37NW right side up joint N67W, 49NE N68W, 54NE joint N08E, 90 N58W, 54NE joint N43W, 62NE N64W, 49NE N23W, 42NE Dipping to north joint N10W, 88NE N53W, 33NE joint N68W, 82NE N65E, 27NW joint N65W, 90 N70E, 26NW joint N40E, 85NW N72W, 23NE N20W, 29NE N74W, 30NE N10E, 53NW N10W, 16NE N08W, 40NE N10E, 51NW N70W, 38NE N20W, 24NE N71E. 40NW N65W, 55NE N23W, 13NE N79W, 32SW N22W, 89NE N58W. 33NE N07E, 47NW N65W. 45NE N52W, 21NE N67W, 14NE contact NE, dip NW N85E, 38NW N40W, 35NE N58W, 48NE N55W, 30NE fold plane N14W, 68NE N42W, 52NE N43W, 87SW EW, 35N N80W, 25NE N41E, 49SE

Table 1. Correspondence of map numbers with field stations

44Hr7/10-7

44Hr7/6-1

44Hr7/7-1

83MR234

83GA231

44Hr7/6-17

44Hr7/6-16

44Hr7/6-15

44Hr7/6-14

44Ca7/11-3

44Hr7/11-3

44Hr7/10-10

44Hr7/10-11

44Hr7/10-2

44Hr7/10-6

44Hr7/10-5

44Hr7/10-3

44Hr7/10-4

44Hr7/7-2

44Ca7/11-2

44Hr7/11-2

44Hr7/11-1

44Hr7/10-1

44Hr7/9-4

44HR7/9-3

Attitude of

bedding and

volcanic flow

44Hr7/9-1

44Hr7/8-4

44Hr7/8-3

44Hr7/8-2

44Hr7/8-11

44Hr7/8-10

44Hr7/8-10B

44Hr7/8-4B

44Hr7/8-5

44Hr7/8-5A

44Hr7/8-5B

44Hr7/8-9

44Hr7/8-5C

44Hr7/8-5D

44Hr7/8-6

44Hr7/8-8

Other structural

data

contact N88E, 37SE

44 Hr 7/8-7

44Hr7/8-1

44Hr7/4-4

44Hr7/4-5

44Hr7/6-6

44Hr7/6-7

44Hr7/6-8

44Hr7/6-9

44Hr7/6-11

44 Ca6/28-6

44Ca6/28-7

44Hr6/28-8

44Ca6/28-8

44Hr6/28-9

44Ca7/4-1a

44Hr7/4-1B

44Hr7/4-1

44Hr7/5-3

82MR309

44Hr7/6-2

44Hr7/6-3

44Hr7/6-4

44Hr7/6-5

44Hr7/6-13

44Hr7/6-12

44Ca7/11-4

44Hr7/10-9

44Hr7/10-8B

\*Year-geologist-month/day-field number or year-geologist-field number; Hr = Joseph Hoare, Ca = W.M. Cady, MR = M.S. Robinson,

Map

Table 3. Structural data.

44Hr7/5-2

Other structural

44Hr7/5-1

44Hr6/29-6

44Ca6/29-6

44Ca6/28-12

44Hr6/28-13

44Ca6/28-13

44Hr6/28-14

83TS88

83TS87

44Hr7/4-9

44Hr7/4-6

44Hr7/6-10

44Hr6/28-6

44Ca6/28-5

44Ca6/29-5

44Hr6/29-5

44Ca6/28-9

44Hr6/28-10

44Ca6/28-10

44Hr6/28-11

44Ca6/28-11

44Hr6/28-12

44Hr7/4-8

44Hr7/4-7

44Hr7/4-2B

44Hr7/4-2

44Hr7/4-3

44Hr7/4-3B

TS = T.N. Smith, GA = G.A. Allegro

Attitude of

bedding and

N60W, 28NE

N69W, 23NE

N25W, 10NE

volcanic flow planes (f)

83 TS86

N35W, 65NE N62W. 38NE N85W, 58SW N55W, 54NE E-W, ?? N68W, 23NE N56W, 45SW N52W, 45NE N80E. 32SE fracture cleavage N80W, 46SW N55W, 09SW cleavage N57E, N40W, 42NE N45W, 43NE 56SE N58W, 47SW N48W, 47NE contact N40E, N70E, 38N cleavage N85E, joint N50W, 80SW sill E-W strike joint N50W, 81SW sill E-W, ?? joint N08E, 53SE joint N48W, 31NE sill E-W, ?? N66W, 19NE

N35W, 20NE N83E, 68SE fold axis N68W, 90 N15W, 37SW N13W, 35SW Table 5. Radiometric age data.

N68W, 37SW

N83W, 39SW

N68W, 38NE

Note: rad = radiogenic;  $\delta$  = standard deviation;  $\lambda_e + \lambda_e' = 0.581 \times 10^{-10} \text{yr}^{-1}$ ;  $\lambda_B = 4.962 \times 10^{-10} \text{yr}^{-1}$ ;  $40 \text{K/K}_{total} = 1.67 \times 10^{-4} \text{ mol/mol}$ Fairbanks, Geophysical Institute Cooperative Geochronology Laboratory

Cady, W.M., Wallace, R.E., Hoare, J.M., and Webber, E.J., 1955, The central Kusko-

tains Quadrangles, Alaska: U.S. Geological Survey Open-file Report 76-606, 92 p.

Platt, J.N., 1955, Photogeologic interpretation of the Taylor Mountains Quadrangle; U.S. Geological Survey unpublished data, scale 1:250,000, 1 sheet.